

Rocio Suarez Ordoñez, Joanna Travis-Roberts, Rasha Abi Hana, Hendree E. Jones.

WHAT WE DID



Surveyed
800
healthcare
providers



across
82
countries




in
3
key
areas

Three key areas in substance use disorder (SUD) treatment:
Factsheet 1 - The availability of SUD treatment centers for women
Factsheet 2 - The prevalence of trauma-informed care
Factsheet 3 - Digital access for women looking for SUD treatment

THE PROJECT AIMS TO IDENTIFY GAPS IN CARE AND PROVIDE RECOMMENDATIONS TO ADDRESS THEM.

Addressing the gap: Global survey finds urgent need for trauma care in women with substance use disorders (SUD)

WHAT WE FOUND




of providers agree that experiences of trauma contribute to the development of Substance Use Disorders

Respondents rated this on a scale from 1 (not at all) to 5 (extremely). Of the total, 68.5% rated this as 5, 26.6% rated it as 4, and the remaining responses were distributed among the lower ratings.



of providers agree that traumatic experiences affect the outcomes of Substance Use Disorder treatment

66% of respondents rated this as 5, 24.8% rated it as 4, with the rest distributed among the lower ratings.



of providers **DO NOT** utilize a standardized tool for screening and / or assessing Trauma and Post-Traumatic Stress Disorder

Among the 30.1% who did use standardized tools, 32% did not specify the tools they employed.




of providers would like to receive training in trauma

WHAT CAN WE DO?

Implement trauma care in all SUD treatment and recovery settings.

HOW?



This is a project by the joint collaboration of

